

NEW FRACTIONAL INEQUALITIES OF OSTROWSKI-GRÜSS TYPE

MEHMET ZEKI SARIKAYA*, HATICE YALDIZ, AND NAGIHAN BASAK

ABSTRACT. In this paper, we improve and further generalize some Ostrowski-Grüss type inequalities for the fractional integrals by using new Montgomery identities.

1. INTRODUCTION AND PRELIMINARY RESULTS

The inequality of Ostrowski gives us an estimate for the deviation of the values of a smooth function from its mean value. More precisely, if $f : [a, b] \rightarrow \mathbb{R}$ is a differentiable function with bounded derivative, then

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[\frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] (b-a) \|f'\|_\infty$$

for every $x \in [a, b]$. Moreover the constant $1/4$ is the best possible.

For some generalizations of this classic fact see the book [12, p.468-484] by Mitrić, Pecarić and Fink. A simple proof of this fact can be done by using the following identity [12]:

If $f : [a, b] \rightarrow \mathbb{R}$ is differentiable on $[a, b]$ with the first derivative f' integrable on $[a, b]$, then Montgomery identity holds:

$$f(x) = \frac{1}{b-a} \int_a^b f(t) dt + \int_a^b P_1(x, t) f'(t) dt,$$

where $P_1(x, t)$ is the Peano kernel defined by

$$P_1(x, t) := \begin{cases} \frac{t-a}{b-a}, & a \leq t < x \\ \frac{t-b}{b-a}, & x \leq t \leq b. \end{cases}$$

This inequality provides an upper bound for the approximation of integral mean of a function f by the functional value $f(x)$ at $x \in [a, b]$. In 2001, , Cheng [1] proved the following Ostrowski-Grüss type integral inequality.

2000 *Mathematics Subject Classification.* 26D15, 41A55, 26D10, 26A33 .

Key words and phrases. Montgomery identity, fractional integral, Ostrowski inequality, Grüss inequality.

*corresponding author.

Theorem 1. *Let $I \subset \mathbb{R}$ be an open interval, $a, b \in I, a < b$. If $f : I \rightarrow \mathbb{R}$ is a differentiable function such that there exist constants $\gamma, \Gamma \in \mathbb{R}$, with $\gamma \leq f'(x) \leq \Gamma$, $x \in [a, b]$. Then have*

$$(1.1) \quad \left| \frac{1}{2}f(x) - \frac{(x-b)f(b) - (x-a)f(a)}{2(b-a)} - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ \leq \frac{(x-a)^2 + (b-x)^2}{8(b-a)} (\Gamma - \gamma), \text{ for all } x \in [a, b].$$

Theorem 1 is a generalization of the following Ostrowski-Grüss type integral inequality, which was firstly by Dragomir and Wang in [2] and further improved by Matic et al. in [6].

Theorem 2. *Let the assumptions of Theorem 1 hold. Then for all $x \in [a, b]$, we have*

$$(1.2) \quad \left| f(x) - \frac{f(b) - f(a)}{b-a} \left(x - \frac{a+b}{2} \right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ \leq \frac{1}{4} (b-a) (\Gamma - \gamma).$$

The above two inequalities are both connections between the Ostrowski inequality [4] and the Grüss inequality [5] and can be applied to bound some special mean and some numerical quadrature rules.

During the past few years many researchers have given considerable attention to the above inequalities and various generalizations, extensions and variants of these inequalities have appeared in the literature, see [1], [2], [6], [23] and the references cited therein. For recent results and generalizations concerning Ostrowski and Grüss inequalities, we refer the reader to the recent papers [1]-[6], [12]-[15], [17]-[23].

The theory of fractional calculus has known an intensive development over the last few decades. It is shown that derivatives and integrals of fractional type provide an adequate mathematical modelling of real objects and processes see ([7], [8], [9], [10], [20], [21]). Therefore, the study of fractional differential equations need more developmental of inequalities of fractional type. The main aim of this work is to develop new weighted Montgomery identity for Riemann-Liouville fractional integrals that will be used to establish new weighted Ostrowski inequalities. Let us begin by introducing this type of inequality.

In [7] and [20], the authors established some inequalities for differentiable mappings which are connected with Ostrowski type inequality by used the Riemann-Liouville fractional integrals, and they used the following lemma to prove their results:

Lemma 1. *Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be differentiable function on I° with $a, b \in I$ ($a < b$) and $f' \in L_1[a, b]$, then*

$$(1.3) \quad f(x) = \frac{\Gamma(\alpha)}{b-a} (b-x)^{1-\alpha} J_a^\alpha f(b) - J_a^{\alpha-1} (P_2(x, b) f(b)) + J_a^\alpha (P_2(x, b) f'(b)), \quad \alpha \geq 1,$$

where $P_2(x, t)$ is the fractional Peano kernel defined by

$$P_2(x, t) = \begin{cases} \frac{t-a}{b-a}(b-x)^{1-\alpha}\Gamma(\alpha), & a \leq t < x \\ \frac{t-b}{b-a}(b-x)^{1-\alpha}\Gamma(\alpha), & x \leq t \leq b. \end{cases}$$

In this article, we use the Riemann-Liouville fractional integrals to establish some new integral inequalities of Ostrowski-Grüss type. From our results, the classical Ostrowski-Grüss type inequalities can be deduced as some special cases.

Firstly, we give some necessary definitions and mathematical preliminaries of fractional calculus theory which are used further in this paper. More details, one can consult [11], [16].

Definition 1. The Riemann-Liouville fractional integral operator of order $\alpha \geq 0$ with $a \geq 0$ is defined as

$$\begin{aligned} J_a^\alpha f(x) &= \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt, \\ J_a^0 f(x) &= f(x). \end{aligned}$$

Recently, many authors have studied a number of inequalities by used the Riemann-Liouville fractional integrals, see ([7], [8], [9], [10], [20], [21]) and the references cited therein.

2. MAIN RESULTS

Lemma 2. Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° with $a, b \in I$ ($a < b$), $\alpha \geq 1$ and $f' \in L_1[a, b]$, then the generalization of the Montgomery identity for fractional integrals holds:

$$\begin{aligned} f(x) &= (\alpha+1)\Gamma(\alpha) \frac{(b-x)^{1-\alpha}}{(b-a)} J_a^\alpha f(b) - J_a^{\alpha-1}(P_2(x, b)f(b)) - \frac{(b-x)^{2-\alpha}}{(b-a)} \Gamma(\alpha) J_a^{\alpha-1} f(b) \\ &\quad - \frac{(b-x)^{1-\alpha}(x-a)}{(b-a)^{2-\alpha}} f(a) + 2J_a^\alpha \left(K_1(x, b) f'(b) \right), \end{aligned} \tag{2.1}$$

where $K_1(x, t)$ is the fractional Peano kernel defined by

$$K_1(x, t) := \begin{cases} \left(t - \frac{a+x}{2} \right) \frac{(b-x)^{1-\alpha}}{b-a} \Gamma(\alpha), & t \in [a, x] \\ \left(t - \frac{b+x}{2} \right) \frac{(b-x)^{1-\alpha}}{b-a} \Gamma(\alpha), & t \in [x, b]. \end{cases} \tag{2.2}$$

Proof. By definition of $K_1(x, t)$, we have

$$\begin{aligned} & J_a^\alpha \left(K_1(x, b) f'(b) \right) \\ &= \frac{1}{\Gamma(\alpha)} \int_a^b (b-t)^{\alpha-1} K_1(x, t) f'(t) dt \\ &= \frac{(b-x)^{1-\alpha}}{b-a} \left[\int_a^x (b-t)^{\alpha-1} \left(t - \frac{a+x}{2} \right) f'(t) dt + \int_x^b (b-t)^{\alpha-1} \left(t - \frac{b+x}{2} \right) f'(t) dt \right] \end{aligned}$$

that can be written as
(2.3)

$$J_a^\alpha \left(K_1(x, b) f'(b) \right) = \frac{1}{2} J_a^\alpha (P_2(x, b) f'(b)) + \frac{(b-x)^{1-\alpha}}{2(b-a)} \int_a^b (b-t)^{\alpha-1} (t-x) f'(t) dt.$$

For term in the right hand side of (2.3) integrating by parts implies that

$$\begin{aligned} & \int_a^b (b-t)^{\alpha-1} (t-x) f'(t) dt \\ (2.4) \quad &= (b-x) \int_a^b (b-t)^{\alpha-1} f'(t) dt - \int_a^b (b-t)^\alpha f'(t) dt \\ &= (x-a)(b-a)^{\alpha-1} f(a) + (b-x) \Gamma(\alpha) J_a^{\alpha-1} f(b) - \Gamma(\alpha+1) J_a^\alpha f(b) \end{aligned}$$

Substituting $J_a^\alpha (P_2(x, b) f'(b))$ in the Lemma 1 and (2.4) in (2.3), we obtain that

$$\begin{aligned} & J_a^\alpha \left(K_1(x, b) f'(b) \right) \\ &= \frac{1}{2} f(x) - (\alpha+1) \Gamma(\alpha) \frac{(b-x)^{1-\alpha}}{2(b-a)} J_a^\alpha f(b) + \frac{1}{2} J_a^{\alpha-1} (P_2(x, b) f(b)) \\ & \quad + \frac{(b-x)^{1-\alpha} (x-a)}{2} (b-a)^{\alpha-2} f(a) + \frac{(b-x)^{2-\alpha}}{2(b-a)} \Gamma(\alpha) J_a^{\alpha-1} f(b). \end{aligned}$$

The proof is completed. $\square$

Remark 1. Letting $\alpha = 1$, formula (2.1) reduces the following identity

$$\frac{1}{2} f(x) = \frac{1}{(b-a)} \int_a^b f(t) dt + \frac{(x-b)f(b) - (x-a)f(a)}{2(b-a)} + \int_a^b K(x, t) f'(t) dt$$

where

$$K(x, t) := \begin{cases} \left(t - \frac{a+x}{2} \right), & t \in [a, x) \\ \left(t - \frac{b+x}{2} \right), & t \in [x, b] \end{cases}$$

which was given by Tong and Guan in [23]. Using the above identity, the authors proved another simple proof of Theorem 1.

Now using the new Montgomery identity for fractional integrals (2.1) and the corresponding fractional Peano kernel (2.2), we derive a new Ostrowski-Grüss type inequality of fractional type.

Theorem 3. *Let f be a differentiable function on $[a, b]$ and $|f'(x)| \leq M$ for any $x \in [a, b]$. Then the following fractional inequality holds:*

$$\begin{aligned}
 & \left| \frac{1}{2} f(x) - (\alpha + 1) \Gamma(\alpha) \frac{(b-x)^{1-\alpha}}{2(b-a)} J_a^\alpha f(b) + \frac{1}{2} J_a^{\alpha-1} (P_2(x, b) f(b)) \right. \\
 & \left. + \frac{(b-x)^{2-\alpha}}{2(b-a)} \Gamma(\alpha) J_a^{\alpha-1} f(b) + \frac{(b-x)^{1-\alpha} (x-a)}{2(b-a)^{2-\alpha}} f(a) \right| \\
 & \leq \frac{M (b-x)^{1-\alpha}}{(b-a)} \left[\frac{(b-a)^\alpha (x-a) + (b-x)^\alpha (a+b-2x)}{2\alpha} \right]
 \end{aligned} \tag{2.5}$$

for $\alpha \geq 1$.

Proof. From Lemma 2, we have

$$\begin{aligned}
 & \left| \frac{1}{2} f(x) - (\alpha + 1) \Gamma(\alpha) \frac{(b-x)^{1-\alpha}}{2(b-a)} J_a^\alpha f(b) + \frac{1}{2} J_a^{\alpha-1} (P_2(x, b) f(b)) \right. \\
 & \left. + \frac{(b-x)^{2-\alpha}}{2(b-a)} \Gamma(\alpha) J_a^{\alpha-1} f(b) + \frac{(b-x)^{1-\alpha} (x-a)}{2(b-a)^{2-\alpha}} f(a) \right| \\
 & = \frac{1}{\Gamma(\alpha)} \left| \int_a^b (b-t)^{\alpha-1} K_1(x, t) f'(t) dt \right|.
 \end{aligned}$$

Taking into account the assumptions on the function f , it yields

$$\begin{aligned}
 & \left| \frac{1}{2} f(x) - (\alpha + 1) \Gamma(\alpha) \frac{(b-x)^{1-\alpha}}{2(b-a)} J_a^\alpha f(b) + \frac{1}{2} J_a^{\alpha-1} (P_2(x, b) f(b)) \right. \\
 & \left. + \frac{(b-x)^{2-\alpha}}{2(b-a)} \Gamma(\alpha) J_a^{\alpha-1} f(b) + \frac{(b-x)^{1-\alpha} (x-a)}{2(b-a)^{2-\alpha}} f(a) \right| \\
 & \leq \frac{M (b-x)^{1-\alpha}}{(b-a)} \left[\int_a^x (b-t)^{\alpha-1} \left| t - \frac{a+x}{2} \right| dt \right. \\
 & \left. + \int_x^b (b-t)^{\alpha-1} \left| t - \frac{b+x}{2} \right| dt \right].
 \end{aligned} \tag{2.6}$$

Noting the left hand side of (2.6) by I then integrating by parts the right hand side of (2.6), we obtain

$$I \leq \frac{M(b-x)^{1-\alpha}}{2(b-a)} \left[\frac{(b-a)^\alpha - (b-x)^\alpha}{\alpha} (x-a) + \frac{(b-x)^{\alpha+1}}{\alpha} \right].$$

Consequently

$$I \leq \frac{M(b-x)^{1-\alpha}}{(b-a)} \left[\frac{(b-a)^\alpha (x-a) + (b-x)^\alpha (a+b-2x)}{2\alpha} \right].$$

This completes the proof. $\square$

Remark 2. Letting $\alpha = 1$, formula (2.5) reduces the following inequality

$$\begin{aligned} & \left| \frac{1}{2}f(x) - \frac{(x-b)f(b) - (x-a)f(a)}{2(b-a)} - \frac{1}{(b-a)} \int_a^b f(t)dt \right| \\ & \leq \frac{M}{(b-a)} \left[\frac{(x-a)^2 + (b-x)^2}{2} \right]. \end{aligned}$$

which connected with Ostrowski-Grüss type integral inequality. If we take $x = \frac{a+b}{2}$ in this inequality, it follows that

$$\begin{aligned} & \left| \frac{1}{2}f\left(\frac{a+b}{2}\right) + \frac{f(b) + f(a)}{4} - \frac{1}{(b-a)} \int_a^b f(t)dt \right| \\ & \leq \frac{M(b-a)}{4}. \end{aligned}$$

REFERENCES

- [1] X. L. Cheng, *Improvement of some Ostrowski-Grüss type inequalities*, Computers Math. Applic, 42 (2001), 109-114.
- [2] S. S. Dragomir and S. Wang, *An inequality of Ostrowski-Grüss type and its applications to the estimation of error bounds for some special means and for some numerical quadrature rules*, Computers Math. Applic, 33(11)(1997), 15-20.
- [3] P. Cerone and S.S. Dragomir, *Trapezoidal type rules from an inequalities point of view*, Handbook of Analytic-Computational Methods in Applied Mathematics, CRC Press N.Y. (2000).
- [4] D. S. Mitrinovic, J. Pecaric and A. M. Fink, *Inequalities for Functions and Their Integrals and Derivatives* Kluwer Academic, Dordrecht, (1994).
- [5] D. S. Mitrinovic, J. Pecaric and A. M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic, Dordrecht, (1993).
- [6] M. Matic, J. Pecaric and N. Ujevic, *Improvement and further generalization of inequalities of Ostrowski-Grüss type*, Computers Math. Appl., 39(3/4), (2000), 161-175.
- [7] G. Anastassiou, M.R. Hooshmandasl, A. Ghasemi and F. Moftakharzadeh, *Montgomery identities for fractional integrals and related fractional inequalities*, J. Inequal. in Pure and Appl. Math, 10(4), 2009, Art. 97, 6 pp.
- [8] S. Belarbi and Z. Dahmani, *On some new fractional integral inequalities*, J. Inequal. in Pure and Appl. Math, 10(3), 2009, Art. 97, 6 pp.
- [9] Z. Dahmani, L. Tabharit and S. Taf, *Some fractional integral inequalities*, Nonlinear Science Letters A, 2(1), 2010, p.155-160.
- [10] Z. Dahmani, L. Tabharit and S. Taf, *New inequalities via Riemann-Liouville fractional integration*, J. Advance Research Sci. Comput., 2(1), 2010, p.40-45.

- [11] R. Gorenflo and F. Mainardi, *Fractional calculus: integral and differentiable equations of fractional order*, Springer Verlag, Wien, 1997, p.223-276.
- [12] D. S. Mitrinovic, J. E. Pecaric and A. M. Fink, *Inequalities involving functions and their integrals and derivatives*, Kluwer Academic Publishers, Dordrecht, 1991.
- [13] S.S. Dragomir and N. S. Barnett, *An Ostrowski type inequality for mappings whose second derivatives are bounded and applications*, RGMIA Research Report Collection, V.U.T., 1(1999), 67-76.
- [14] S.S. Dragomir, *An Ostrowski type inequality for convex functions*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. 16 (2005), 12-25.
- [15] Z. Liu, *Some companions of an Ostrowski type inequality and application*, J. Inequal. in Pure and Appl. Math, 10(2), 2009, Art. 52, 12 pp.
- [16] S. G. Samko, A. A Kilbas, O. I. Marichev, *Fractional Integrals and Derivatives Theory and Application*, Gordon and Breach Science, New York, 1993.
- [17] M. Z. Sarikaya, *On the Ostrowski type integral inequality*, Acta Math. Univ. Comenianae, Vol. LXXIX, 1(2010), pp. 129-134.
- [18] M. Z. Sarikaya, *On the Ostrowski type integral inequality for double integrals*, Demonstratio Mathematica, accepted.
- [19] M. Z. Sarikaya and H. Ogunmez, *On the weighted Ostrowski type integral inequality for double integrals*, The Arabian Journal for Science and Engineering (AJSE)-Mathematics, (2011) 36: 1153-1160
- [20] M.Z. Sarikaya and H. Ogunmez, *On new inequalities via Riemann-Liouville fractional integration*, arXiv:1005.1167v1, submitted.
- [21] M.Z. Sarikaya, E. Set, H. Yaldiz and N., Basak, *Hermite -Hadamard's inequalities for fractional integrals and related fractional inequalities*, Mathematical and Computer Modelling, DOI:10.1016/j.mcm.2011.12.048.
- [22] A. Rafiq and F. Ahmad, *Another weighted Ostrowski-Grüss type inequality for twice differentiable mappings*, Kragujevac Journal of Mathematics, 31 (2008), 43-51.
- [23] F. Tong and L. Guan, *A simple proof of the generalized Ostrowski-Grüss type integral inequality*, Int. Jour. of Math. Analysis, 2(18), (2008), 889-892.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, DÜZCE, TURKEY

E-mail address: sarikayamz@gmail.com

E-mail address: yaldizhatice@gmail.com

E-mail address: nagihan.basak@hotmail.com